

Florigen and the Photoperiodic Control of Flowering in Rice

Hiroyuki Tsuji · Shojiro Tamaki · Reina Komiya ·
Ko Shimamoto

Received: 19 May 2008 / Accepted: 25 June 2008 / Published online: 15 August 2008
© Springer Science + Business Media, LLC 2008

Abstract Flowering time is a key trait for geographical and seasonal adaptation of plants and is an important consideration for rice breeders. Recently identified genetic factors provide new insights into this complex trait. The list of genes involved in flowering and their functions tells us that the molecular basis of day-length measurement includes both of the evolution of unique factors and the regulatory adaptation of conserved factors in rice. This information helped identify rice florigen, a mobile flowering signal. Our current view of flowering time regulation incorporates the presence of complex layers of gene networks integrated with the synthesis of florigen protein and its subsequent transport and perception.

Keywords Florigen · Flowering time · Genetic pathways

Introduction

Flowering time, often referred to as heading date in cereal crops, is a key agronomical determinant for adaptation to specific cropping locations and growing seasons for current varieties of cultivated rice. Developing early-flowering or photoperiod-insensitive cultivars has been a major objective of rice breeding for several decades. The molecular and developmental determinants of flowering time have thus also particularly important genetic targets for domestication or for breeding new varieties of rice. Flowering time is

controlled by many genes, which are expressed or suppressed in close interaction with environmental factors such as day length and temperature. Progress in the molecular genetics has provided a clearer understanding of several pivotal mechanisms regulating flowering time determination in rice, and these studies have been summarized from genetic, molecular biological, or comparative biological perspectives [17, 24, 25, 43, 83]. In this review, we discuss the current understanding of rice flowering, considering the recent finding of a strong candidate for the long sought-after mobile flowering stimulus, florigen [74].

Hd3a protein as the rice florigen

Heading date 3a (Hd3a) was first detected as a quantitative trait locus (QTL), which promotes flowering of rice under short-day (SD) conditions [55, 80]. *Hd3a* encodes an ortholog of *Arabidopsis FLOWERING LOCUS T (FT)* [38]. Overexpression of Hd3a protein with the constitutive promoter [38] or vascular-specific promoters [74] results in an early-flowering phenotype, and suppression of *Hd3a* with RNA interference (RNAi) delays flowering [39]. *Hd3a* is a member of a large gene family consisting of at least 13 genes in rice genome [6] and at least two paralogs, *Rice FLOWERING LOCUS T1 (RFT1)* and *FT-Like (FTL)* promote flowering [28, 39]. Hd3a/FT proteins are about 22 kDa in size, and their overall structures are similar to mammalian phosphatidyl ethanolamine-binding proteins or Raf-kinase inhibitor proteins (RKIP) [3], although it is not known whether or not Hd3a/FT binds to phosphatidyl ethanolamine or inhibits kinase.

Recent progress in the molecular genetics of *Arabidopsis* indicates that *FT* has a strong flowering promotion effect downstream of the known photoperiodic flowering pathway

H. Tsuji · S. Tamaki · R. Komiya · K. Shimamoto (✉)
Laboratory of Plant Molecular Genetics,
Nara Institute of Science and Technology (NAIST),
8916-5 Takayama,
Ikoma, Nara 630-0101, Japan
e-mail: simamoto@bs.naist.jp

[17, 21, 36]. *Arabidopsis* flowering is promoted by long day (LD), whereas rice flowering is promoted by SD. *FT* is a major floral activator that is expressed in the vascular tissues of leaves under inductive LD in *Arabidopsis* [32, 37, 71]. *FT* protein interacts with the bZIP transcription factor FLOWERING LOCUS D (FD), which is only expressed in the shoot apical meristem (SAM), and this interaction seems to be necessary for full activation of *FT* floral promotion [1, 76]. The anatomical separation of expression sites implies that *FT* product is a primary candidate for florigen. If *FT* encodes the florigen, it could act as a mobile agent either via protein or messenger RNA (mRNA), either of which could move to the SAM to interact with FD. Therefore, a major focus was whether transcripts or protein account for the mobile florigenic activity [85].

The question was answered with the introduction of rice as a paradigm for exploring the nature of florigen (Fig. 1) [74]. First, the precise sites of *Hd3a* transcription and mRNA accumulation were determined. The promoter activity of *Hd3a* was detected in transgenic plants expressing β -glucuronidase (GUS) under the control of the *Hd3a* promoter in the vascular cells of leaf blades. No GUS activity was detected in the SAM. This is consistent with the tissue specificity of *FT* expression in *Arabidopsis*, although expression is induced by LD for *FT* and by SD for *Hd3a* [32, 37]. *Hd3a* mRNA accumulates in leaf blades but is present at very low levels in leaf sheaths and is four orders of magnitude lower in the shoot apex than in leaf blades under SD conditions. Therefore, it is unlikely that *Hd3a* mRNA moves from leaves to the SAM in any significant amount. *Hd3a* expression is thus

limited to the vascular tissues of leaf blades under SD conditions.

Secondly, the tissue localization of Hd3a protein was examined using transgenic rice plants expressing Hd3a-GFP fusion protein driven by the Hd3a promoter. Hd3a-GFP was present in vascular tissue from the leaf blade to the upper part of the stem, the region just below the meristem where nodes are present, and in the inner cone-like region of the SAM. Hd3a-GFP protein is thus synthesized in the vascular tissues of leaf blades and transported through the phloem. Hd3a protein is then unloaded from the end of the vascular tissue and enters the SAM through cells just beneath the SAM [74]. Essentially, the same localization of Hd3a-GFP was observed when other vascular-specific promoters were used to express Hd3a-GFP, confirming that Hd3a-GFP is translocated from leaf and stem vasculature to the SAM. These observations indicate that Hd3a protein is a mobile flowering signal in rice (Fig. 1). The ability of Hd3a/FT protein to move long distances is also suggested from a study of the rice phloem sap proteome [4]. Rice phloem sap contains a member of Hd3a/FT protein family, FT-LIKE12, along with RKIP-like proteins RICE CENTRORADIALIS3 (RCN3) and MOTHER OF FT1. Hd3a itself was not detected, probably because photoperiod conditions and timing of sampling was not suitable for detection of Hd3a. Hd3a/FT proteins, however, have been detected in the phloem sap of other plant species. CmFTL2, a functional ortholog of Hd3a/FT in *Cucurbita maxima*, has also been found in vascular sap and can move through graft unions to promote flowering [47]. Phloem sap obtained from the inflorescence




Fig. 1 Model of Hd3a/FT action in rice and *Arabidopsis*. Hd3a/FT protein is synthesized in the vascular tissue of the leaf and transported through phloem tissue. Hd3a/FT protein is unloaded from the end of the vascular tissue and enters into the SAM through cells just

below the SAM. Several lines of evidence suggest that Hd3a/FT protein interacts with the bZIP transcription factor FD, which directly binds promoter region of floral meristem identity gene *AP1*. Hd3a/FT is a strong candidate for florigen.

stems of *Brassica napus* also contained FT protein [14]. These observations provide additional evidence that Hd3a/FT proteins are florigens. Evidence from *Arabidopsis* also supports the florigen activity of FT protein. Phloem-expressed FT-GFP and Myc-tagged FT move to the apex and cause early flowering, but FT protein with nuclear localization signal or 3xYFP do not, probably because of immobilization [8, 29, 52].

RFT1 also promotes flowering in rice

The *RFT1* gene lies adjacent to *Hd3a*, separated by only 11.5 kb on chromosome 6 of rice [26]. *Hd3a* and *RFT1* sequences are very similar, and they were likely produced by a recent tandem duplication event [6]. The corresponding chromosomal regions of the *Hd3a/RFT1* locus in maize or wheat contain only a single *Hd3a* ortholog [10, 82]. *RFT1* tissue-specific expression is similar to that of *Hd3a* in that it is expressed in vascular tissues of leaf blades in response to SD. Thus, *RFT1* could possibly function redundantly with *Hd3a* in promoting rice flowering as another florigen. The interesting involvement of *RFT1* was revealed from a detailed analysis of rice plants with RNAi-suppressed *Hd3a* expression [39]. Flowering of *Hd3a* RNAi plants was delayed by more than 30 days under SD conditions. Although flowering of *RFT1* RNAi plants was normal, suggesting that *RFT1* does not on its own promote flowering under SD conditions, double RNAi plants in which both *Hd3a* and *RFT1* are suppressed did not flower up to 300 days after germination. Both *Hd3a* and *RFT1* are thus involved in the normal regulation of flowering in rice. *RFT1* expression is normally low in wild-type rice plants but is up-regulated in *Hd3a* RNAi plants and is correlated with reproductive transition of the SAM. This up-regulation is also correlated with an increase of histone-3-acetylation of nucleosomes around the transcription initiation site of *RFT1*. These results suggest that *RFT1* expression increases in the absence of Hd3a and that *RFT1* complements *Hd3a* function [39]. RFT1 thus serves as another flowering promotion activity in rice under certain conditions.

Hd3a/FT signaling in the shoot apical meristem

The identification of Hd3a/FT as florigen opens the door to studying the molecular mechanism(s) of Hd3a/FT signaling in the SAM. The current understanding of events downstream of Hd3a/FT activity focuses on FD, MADS-box transcription factors, and CENTRORADIALIS (CEN) / TERMINAL FLOWER1 (TFL1). Corresponding genes of the proteins discussed below are summarized in Table 1.

FD

In the SAM, Hd3a may interact with a bZIP transcription factor orthologous to *Arabidopsis* FD to induce phase transition from vegetative to reproductive states. In *Arabidopsis*, *FD* is expressed on the flanks of the SAM where flower primordia are initiated. *FD* is expressed during the vegetative phase so that it is present when FT arrives. In the presence of FT, FD induces the floral meristem identity gene *APETALA1* (*AP1*) and its paralog *FRUITFULL* (*FUL*) [2, 76]. At present, there are no reports describing FD function in rice, whereas the maize late flowering mutant *delayed flowering 1* (*dlf1*) provides a hint about FD function in rice and other cereals [58]. The *DLF1* gene encodes an ortholog of *FD*. Loss-of-function mutants of maize *DLF1* (*dlf1*) have a late flowering phenotype with some abnormal tassel branches. The most abundant *DLF1* expression is found in the SAM, and it peaks during vegetative-to-reproductive phase transition, decreasing gradually during inflorescence development. *DLF1* has also been shown to interact with maize Hd3a/FT. The maize genome contains at least 15 members of the *Hd3a/FT* gene family. Two of them, *Zea mays* *CENTRORADIALIS 8* (*ZCN8*) and *ZCN14*, interact with *DLF1* in yeast two-hybrid assays [10]. *ZCN8* transcripts are present in leaf blades before floral transition, increases just before floral transition, and stays high afterward. Because of its expression pattern and interaction with *DLF1*, *ZCN8* should be considered as a florigen candidate in maize. *ZCN15*, an ortholog of rice *Hd3a*, which resides in the syntenic chromosome region of rice *Hd3a*, is weakly expressed in leaf blades after phase transition and relatively highly expressed in the developing kernel. The activation of *AP1* by Hd3a/FT and FD is operative in cereal crops. The *dlf1* mutation of maize delayed the expression of maize *AP1/FUL* MADS-box gene *ZMM4*, an ortholog of *OsMADS14/RAP1B*, in the SAM [11], and TaFDL2, a wheat FD homolog, can interact with wheat FT protein *VERNALIZATION3* (*VRN3*)/*TaFT* and directly binds the promoter region of wheat *AP1* homologous gene in vitro [45].

There are at least seven orthologs of *FD* in the rice genome [60]. *OsZIP24* mRNA is relatively abundant in the SAM, and *OsZIP54* mRNA is distributed throughout the rice plant, including the SAM, whereas transcript accumulation of other FD-like bZIPs is below the microarray detection limit [60]. It is yet to be determined if there is some unresolved functional differentiation among these FD-like proteins of rice.

MADS-box transcription factors

The gene network, which acts downstream of FT in the SAM, involves *MADS-box* genes in *Arabidopsis*. Both FT

Table 1 Genes that are Expected to be Involved in Hd3a/FT Signaling in SAM

| Category | Gene name | Accession no. | Organism | Domain | Function | Reference |
|--------------------------------------|---|-----------------------|--------------------|----------------------|----------------------------------|------------------------|
| Hd3a/FT | <i>Hd3a</i> | Os06g0157700 | Rice | Raf-kinase inhibitor | Florigen; promotion of flowering | [38] |
| | <i>RFT</i> | Os06g0157500 | Rice | | Promotion of flowering | [28] |
| | <i>FTL</i> | Os01g0218500 | Rice | | Promotion of flowering | [28] |
| | <i>ZCN8</i> | EU241924 | Maize | | n.d. | [11] |
| | <i>ZCN15</i> | EU241930 | Maize | | n.d. | [11] |
| | <i>VRN3/TaFT</i> | DQ890165 | Wheat | | Promotion of flowering | [82] |
| | <i>FT</i> | At1g65480 | <i>Arabidopsis</i> | | Florigen; Promotion of flowering | [32, 37] |
| CEN/TFL1 | <i>TSF</i> | AT4g20370 | <i>Arabidopsis</i> | | Promotion of flowering | [78] |
| | <i>RCN1</i> | Os11g0152500 | Rice | Raf-kinase inhibitor | Inhibition of flowering | [59] |
| | <i>RCN2</i> | Os02g0531600 | Rice | | Inhibition of flowering | [59] |
| | <i>RCN3</i> | Os12g0152000 | Rice | | Inhibition of flowering | [59, 87] |
| | <i>OsCEN3</i> | Os04g0411400 | Rice | | n.d. | [87] |
| | <i>TFL1</i> | At5g03840 | <i>Arabidopsis</i> | | Inhibition of flowering | [5] |
| FD | <i>OsbZIP24</i> | Os02g0833600 | Rice | bZIP | n.d. | [60] |
| | <i>OsbZIP27</i> | Os03g0306700 | Rice | | n.d. | [60] |
| | <i>OsbZIP54</i> | Os06g0719500 | Rice | | n.d. | [60] |
| | <i>OsbZIP55</i> | Os06g0720900 | Rice | | n.d. | [60] |
| | <i>OsbZIP56</i> | Os06g0724000 | Rice | | n.d. | [60] |
| | <i>OsbZIP69</i> | Os08g0549600 | Rice | | n.d. | [60] |
| | <i>OsbZIP77</i> | Os09g0540800 | Rice | | n.d. | [60] |
| | <i>DLF1</i> | EF093789 | Maize | | Promotion of flowering | [58] |
| | <i>TaFDL2</i> | EU307112 | Wheat | | n.d. | [45] |
| | <i>FD</i> | At4g35900 | <i>Arabidopsis</i> | | Promotion of flowering | [1, 76] |
| | MADS-box transcription factor (FUL/AP1/CAL) | <i>OsMADS14/RAP1B</i> | Os03g0752800 | Rice | MADS-box | Promotion of flowering |
| <i>OsMADS15/RAP1A</i> | | Os07g0108900 | Rice | | n.d. | [41] |
| <i>ZMM4</i> | | AJ430641 | Maize | | Promotion of flowering | [11] |
| <i>WAP1/VRN1</i> | | AB007504/AY188331 | Wheat | | Promotion of flowering | [56, 81] |
| <i>FUL</i> | | At5g60910 | <i>Arabidopsis</i> | | Promotion of flowering | [15] |
| <i>API</i> | | At1g69120 | <i>Arabidopsis</i> | | n.d. | [51] |
| <i>CAL</i> | | At1g26310 | <i>Arabidopsis</i> | | n.d. | [33] |
| MADS-box transcription factor (SOC1) | <i>OsMADS50/OsSOC1</i> | Os03g0122600 | Rice | MADS-box | Promotion of flowering | [44] |
| | <i>SOC1</i> | At2g45660 | <i>Arabidopsis</i> | | Promotion of flowering | [65] |

n.d. Not functionally determined yet

and FD, together with LEAFY, are required for inducing expression of *API* and *FUL* in the SAM [2, 53, 75]. FT-dependent gene regulation is also observed in *Arabidopsis* leaves. Expression of *FUL* and *SEPALATA3* was induced in leaves overexpressing *FT* and was reduced in *ft* and *fd* mutants [75]. Thus, leaves and the SAM may share similar regulatory mechanisms. In rice, the expression of *OsMADS14/RAP1B*, an ortholog of *API*, and *OsMADS15/RAP1A* are up-regulated by *Hd3a* in leaf blades under SD conditions [39], suggesting that the same regulatory network is involved in the SAM of rice. *OsMADS14/RAP1B* is expressed in the SAM during the reproductive phase [13, 41], and its ectopic expression strongly promotes rice flowering [31]. Interestingly, a mutant of the *OsMADS14/RAP1B* ortholog in einkorn wheat (*Triticum monococcum*), *maintained vegetative phase (mvp)*, cannot show transition

from vegetative to reproductive phases [67]. The *mvp* phenotype is caused by a deletion of the promoter and coding region of *Wheat API (WAP1)/VERNALIZATION1 (VRN1)*, a gene that lies within the wheat chromosome region corresponding to the region containing *OsMADS14/RAP1B* in rice [81]. Reduced levels of *VRN3/TaFT*, an ortholog of rice *Hd3a*, are associated with lower levels of *VRN1* transcripts, consistent with the observation in rice [82]. Although *API* in *Arabidopsis* is unlikely to be the FT target in the SAM for flowering control because *ap1* mutants are not late flowering [62], *API* MADS-box proteins are likely to function as important mediators of Hd3a/FT activity in flowering of cereals. *OsMADS14/RAP1B* is also an activator of *Hd3a* expression in leaves [34, 43]. Consistent with this, reduced levels of *VRN1* are also correlated with reduced levels of *VRN3* in barley [20].

Suppressor of Overexpression of Constans1 (SOC1) in *Arabidopsis* is a MADS-box protein, and a recent genetic study suggests that FT mediates the activation of *SOC1* downstream of *CONSTANS* (CO) [44]. SOC1 directly interacts with AGAMOUS-LIKE24 (AGL24) MADS-box protein in the shoot apex, and direct binding of the SOC1-AGL24 complex to the promoters of both *SOC1* and *AGL24* upregulates their expression [48]. On the other hand, the regulatory hierarchy and the site of action seem to be different in rice; a T-DNA insertion mutation of *OsMADS50/OsSOC1* results in a reduction of *Hd3a* transcription in leaves, whereas RNAi suppression of *Hd3a* or of both *Hd3a* and *RFT1* does not result in any change in *OsMADS50/OsSOC1* expression. Thus, in rice, *OsMADS50/OsSOC1* acts upstream of *Hd3a* expression in leaf blades [39, 44].

CEN/TFL1

CENTRORADIALIS (CEN)/TERMINAL FLOWER1 (TFL1) proteins provide interesting insights into Hd3a/FT signaling in the SAM. *CEN/TFL1* are homologs of *Hd3a/FT* and encode RKIP proteins, but *CEN/TFL1* proteins repress flowering. *Arabidopsis TFL1* mRNA is limited to the central cells of the inflorescence shoot meristem, and the TFL1 protein migrates to outer cells [7]. Repression of flowering has been thoroughly investigated using *Arabidopsis FT* and TFL1 [3, 16]. Proteins carrying replacements of individual residues or specific regions of FT and TFL1 were overexpressed in a *ft tfl1* double mutant to map the residues or regions that confer specific activity on the two proteins. The crystal structures of FT and TFL1 were also compared to map the region of specificity. These analyses suggest that His88 in TFL1 and the corresponding Tyr85 in FT are important for the opposite activities of FT and TFL1 [16] and that the region responsible for the difference is localized in a 14 amino acid segment in the external loop at the C-termini of the proteins [3].

There are at least four *CEN/TFL1*-like genes in the rice genome [6]. *RICE CENTRORADIALS1 (RCN1)*, *RCN2*, and *RCN3/Oscen1* are able to delay flowering and affect panicle architecture when overexpressed [59, 87]. Interestingly, RCN3 protein can be detected in the phloem sap of rice, suggesting that this protein signals to communicate between distant organs, much like Hd3a [4]. An attractive hypothesis to explain the difference between Hd3a/FT and *CEN/TFL1* regulatory activity is that the external loops bring different partners to Hd3a/FT or *CEN/TFL1*, thus defining the function of the complex. Screening for interacting proteins using specific external loop regions may provide further insights into Hd3a/FT signaling for floral promotion.

Molecular-genetic pathways of the photoperiodic flowering in rice

From an understanding of the molecular nature of Hd3a/FT proteins, it can be inferred that the timing of rice flowering is basically determined by the expression levels of the two essential flowering promotion genes, *Hd3a* and *RFT1* in leaf blades, and by the competency of the SAM for Hd3a/FT activity. Recent molecular genetic studies provide important insights into the day-length regulation of *Hd3a* expression. The crucial aspect of this regulation is SD induction and LD repression of *Hd3a* (Fig. 2).


Short-day promotion of Hd3a expression

Under SD conditions, Hd3a expression is induced gradually and peaks at about 30 days before flowering, about the time floral transition occurs [39]. The diurnal peak of *Hd3a* promoter activity and mRNA accumulation coincide at dawn, suggesting the major point of *Hd3a* regulation is transcriptional [23, 38, 74]. The most important factors for this *Hd3a* expression are *Heading date1 (Hd1)* and *Early heading date1 (Ehd1, see below)* [24]. *Hd1* was first detected as the major photoperiod-sensitivity QTL, and it encodes a B-box zinc finger protein with a C-terminal CCT (CONSTANS, CONSTANS-LIKE and TIMING OF CAB EXPRESSION1) domain [84]. Under SD conditions, loss-of-function alleles of *Hd1* delay flowering and reduce *Hd3a* mRNA accumulation [23, 28, 38, 84]. Detailed phylogenetic analyses showed that *Hd1* is the sole ortholog of *Arabidopsis CONSTANS (CO)*, which is also an important activator of *FT* [26]. *Arabidopsis CO* is expressed in leaf phloem tissues [71], and CO protein seems to directly regulate *FT* and its paralog *TWIN SISTER OF FT (TSF)* because both *FT* and *TSF* are up-regulated by dexamethasone-induced activation of CO in the presence of the translation inhibitor cycloheximide [65, 78]. Whether Hd1 directly regulates *Hd3a* transcription is not clear, but differences in the temporal patterns of *Hd1* and *Hd3a* mRNA accumulation suggest the involvement of a somewhat more complex regulatory mechanism. *Hd1* mRNA levels peak at midnight, but *Hd3a* mRNA levels peak about 4–6 h after *Hd1* [19, 23, 28].

In addition to transcriptional regulation, Hd1 activity may be regulated by protein stability because *Arabidopsis CO* is subject to the regulation at the protein accumulation level. CO protein is destabilized in darkness by CONSTITUTIVELY PHOTOMORPHOGENIC1, which is counteracted by the blue-light photoreceptor CRYPTOCHROME. A PhyB-mediated signal also destabilizes CO at dawn [30, 42, 49].

Phytochromes also affect the expression of *Hd3a* under SD conditions. *Hd3a* expression under SD conditions is

Fig. 2 Regulatory network of rice flowering time. Under short-day conditions, *Hd3a* is mainly up-regulated by Hd1 and Ehd1. *Hd1* is up-regulated by OsGI, and *Ehd1* is up-regulated by OsMADS51, whose expression is up-regulated by OsGI. *OsGI* is regulated by the circadian clock. Under long-day conditions, Hd1 function is converted to repress *Hd3a* in response to a signal from phytochrome. *Ehd1* is repressed by Ghd7, a regulatory component that is up-regulated under long-day conditions, resulting in *Hd3a* repression. SD or LD in parentheses indicate short-day- or long-day-specific regulation, respectively.



significantly up-regulated in an early-flowering *photoperiodic sensitivity 5 (se5)* mutant, which has a mutation in the gene encoding heme oxygenase, which is required for phytochrome chromophore biogenesis [27, 28]. Flowering under SD conditions is also promoted in *phyB* mutants that

have loss-of-function alleles in one of the three rice phytochromes [73]. *Hd1* expression is not affected by *se5* or *phyB* mutations [23, 28]; thus, phytochrome represses *Hd3a* expression downstream or independent of *Hd1* expression under SD conditions. These phytochrome

effects may be reminiscent of the activator-to-repressor conversion of a subset of Hd1 proteins that is observed under LD (see below; Izawa et al. [28]).

The primary upstream regulator of *Hd1* expression is *OsGI*, an ortholog of *GIGANTEA* (*GI*) in *Arabidopsis* [18]. *OsGI* is a large protein that is present in both the nucleus and cytosol of rice cells [1]. Suppression of *OsGI* expression by RNAi or antisense expression caused late flowering and reduced *Hd1* mRNA accumulation under SD conditions [19, 34]. *Arabidopsis* *GI* plays important roles in *CO* expression through the degradation of a repressor of *CO* transcription [22, 66]. Thus, it is possible that *OsGI* is part of a similar mechanism in rice, even though the peak in *OsGI* expression at dusk is quite different from the *Hd1* expression peak at midnight, suggesting the intervention of an unknown mechanism, which caused the difference in timing of expression.

OsGI expression shows circadian daily oscillation, implying upstream regulation by the circadian clock [18]. Circadian clock mutants of *Arabidopsis* have aberrant *GI* expression [54, 61], and *GI* itself is also considered to be a clock-associated protein [35, 50, 86].

Another major *Hd3a* activator, *Ehd1*, was identified from a QTL of a cross between the cultivar T65 and Nipponbare [12]. *Ehd1* encodes a B-type response regulator, and a functional mutation is found in the GARP (maize GOLDEN2, the ARR B-class proteins from *Arabidopsis*, and *Chlamydomonas* *Psr1*) domain, which decreased affinity to the target DNA sequence. Mutations or RNAi suppression of *Ehd1* decreased *Hd3a* expression under SD conditions, indicating that *Ehd1* is the upstream regulator of *Hd3a* expression [12, 34]. *Ehd1* is preferentially expressed under SD conditions, peaking twice, before and after dawn, and *Hd3a* expression is essentially the same as *Ehd1* in the absence of functional *Hd1* [12].

Ehd1 is regulated by at least two factors, *OsMADS51* and *Ghd7* (for grain number, plant height, and heading date 7) [34, 77]. *OsMADS51* acts as the activator of *Ehd1* expression under both SD and LD, while *Ghd7* mainly acts as a suppressor of *Ehd1* under LD (see below). An *osmads51* T-DNA insertion mutation reduced *Ehd1* expression under both SD and LD. *OsMADS51* expression is also up-regulated by *OsGI* because *OsGI* antisense repression reduced *OsMADS51* and *Ehd1* expression. *OsGI* and *OsMADS51* show similar daily expression oscillation, peaking at dusk in SD.

Long-day suppression of *Hd3a* expression

Under LD conditions, *Hd3a* expression is very low in any developmental stage [39]. Diurnal expression is also lower than under inductive SD conditions [38]. Hd1 also has an important role in this suppression, although it works as

activator of *Hd3a* expression under SD. Before the molecular cloning of *Hd1*, it was demonstrated that near-isogenic lines containing *hd1* mutant alleles exhibited not only delayed flowering under SD conditions but also early flowering under LD conditions, indicating that Hd1 could repress flowering under LD and that Hd1 function is modified depending on day length [46]. Phytochrome signaling affects this day-length-dependent conversion of Hd1 activity because this conversion is not observed in the phytochrome-deficient mutant; flowering of the double mutant *se5 hd1* (which lacks both phytochrome and functional *Hd1*) was slightly later than that of the *se5* single mutant (which lacks only phytochrome) [28]. Consistent with this observation, *Hd3a* mRNA accumulation is lower in *se5 hd1* than in *se5* mutant lines. The involvement of functional modification of Hd1 by day length is further supported by the direct manipulation of *OsGI* expression and the resulting change in *Hd1* and *Hd3a* expression levels [19]. Overexpression of *OsGI* delays flowering under both SD and LD, indicating that *OsGI* can act as a suppressor of flowering signals when it is constitutively expressed. In *OsGI* overexpressing plants, *Hd1* mRNA levels increase under both SD and LD, whereas *Hd3a* mRNA levels are negatively correlated with *Hd1* mRNA, indicating that *Hd1* can act as a suppressor of *Hd3a* when it is highly induced by *OsGI* constitutive expression. *OsGI* overexpression could also be expected to increase *Hd3a* expression through *Ehd1* up-regulation by the *OsGI*-*OsMADS51* pathway (see above, Kim et al. [34]), but this effect was not observed, probably because of strong Hd1 suppressive activity. A comparison of daily temporal expression patterns of *Hd1* and *Hd3a* indicate that higher expression levels of *Hd1* during light periods might suppress *Hd3a*. Under SD conditions, wild-type *Hd1* expression starts to increase at the beginning of the dark period, thus keeping Hd1 as an activator. *OsGI* overexpressors, however, express a significant level of *Hd1* during light periods, thus converting Hd1 into a repressor. Under LD, wild-type *Hd1* expression starts to increase in the light, resulting in the conversion of Hd1 into a suppressor [19]. Thus, the external coincidence model could be applied to rice: Circadian clock-regulated *Hd1* expression and an external light signal mediated by phytochrome generate the specific response to day length [25].

The repressor function of Hd1 is not observed in connection with *Arabidopsis* *CO*, indicating that there is a different mechanism that provides this function. The photoperiod-sensitivity QTL *Hd2* may provide information for understanding this mechanism. *Hd2* is detected as a QTL in crosses between *japonica* cultivar Nipponbare and *indica* cultivar Kasalath [80]. A functional allele of *Hd1* is required for Nipponbare allele of *Hd2* to delay flowering in response to LD [46]. In addition, this *Hd2* allele is required

for *Hd6*, another photoperiod-sensitivity QTL, to delay flowering under LD [79]. Map-based cloning of this QTL revealed that it encodes an α -subunit of casein kinase II (CK2 α) [72]. Plant CK2 is composed of two catalytic (α) subunits and two regulatory (β) subunits. CK2 α is mainly localized in nucleus in maize [64], and several nuclear proteins are identified as substrates of CK2 α in maize and barley [40, 68], suggesting that rice *Hd6* also functions in nucleus. In *Arabidopsis*, CK2 activity affects the timing of floral transition through phosphorylation of CIRCADIAN CLOCK ASSOCIATED1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY) proteins, which are the components of central oscillators of plant circadian clock [9, 63, 69, 70]. Thus, the functional interaction between rice LHY and *Hd6* will be an interesting issue. Rice genome contains a single ortholog of *CCA1* and *LHY*, *OsLHY* [26]. When overexpressed in *Arabidopsis*, both *OsLHY* and *Arabidopsis* *CCA1* delay flowering. However, *OsLHY* overexpression showed only subtle effect on the rhythmic expression of *PRR1* compared with the dramatic effect of *CCA1* overexpression, suggesting functional difference between them [57].

Ehd1 expression is suppressed under LD, which also results in reduced *Hd3a* expression. *Ghd7* is responsible for the suppression of *Ehd1* [77]. *Ghd7* protein contains a CCT domain with about 60% identity to *Hd1*, but *Ghd7* lacks the zinc finger motif, which is present in *Hd1*. *Ghd7* expression is significantly higher under LD in leaf vascular tissue, and its diurnal expression peaks at dusk. *Ghd7* expression does not affect *Hd1* expression, but it strongly suppresses *Ehd1*.

Night-break suppression of *Hd3a* expression

A short light exposure during the dark period can significantly delay flowering of SD plants. This night-break effect has long been discussed in the context of day-length measurement in SD plants, but detailed understanding of the molecular mechanism of night break has been very limited. Like other SD plants, rice shows significant response to night break [23]. Night break suppresses *Hd3a* at the transcription level without affecting the expression patterns of *Hd1* and *OsGI*. The effect of illumination varies depending on its timing within the dark period. The night-break effect is most significant in the middle of the dark period. This night-break effect dependence on timing is correlated with the expression pattern of *Hd1*. Because *Hd3a* expression is mainly governed by the activity of *Hd1* and *Ehd1*, the night-break signal may affect both of the activities, which repress *Hd3a* transcription. The night-break effect on *Hd3a* expression was abolished by mutations of *phyB*, thus the night-break signal is mediated by *phyB*, which probably acts via the activities of *Hd1* and *Ehd1* [23].

Perspectives

Research on flowering time regulation in rice expands the field of mobile molecular signaling in plants. Florigen is also referred to as a flowering hormone because of its mobile nature and the universality of its flowering promotion activity. Using the analogy of the canonical plant hormone studies, the next step in florigen research would be understanding *Hd3a*/FT synthesis, transport, identification of receptors, and subsequent cellular signaling. Virtually nothing is known about *Hd3a* long-distance trafficking. How *Hd3a* in companion cells is transferred to sieve element systems, how the direction of movement in phloem is determined, and how *Hd3a* targets the SAM after unloading from the upper end of the phloem are all open questions. Flowering time research has also identified many important genes as resources for breeding. The genetic basis of natural variation or artificial selection in flowering time genes, including genes that are or were lost through the process of domestication, should contribute not only to a further understanding of the mechanism of flowering but also to future breeding applications.

Acknowledgements We thank the members of our laboratory for helpful comments or other assistance. The work described is supported by Grants-in-Aid for Scientific Research on Priority Areas (grant 10182102 and 19090013) of the Ministry of Education, Culture, Sports, Science, and Technology of Japan (to K.S.).

References

1. Abe M, Fujiwara M, Kurotani K, Yokoi S, Shimamoto K. Identification of dynamin as an interactor of rice GIGANTEA by tandem affinity purification (TAP). *Plant Cell Physiol.* 2008;49:420–32.
2. Abe M, Kobayashi Y, Yamamoto S, Daimon Y, Yamaguchi A, Ikeda Y, Ichinoki H, Notaguchi M, Goto K, Araki T. FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* 2005;309:1052–6.
3. Ahn JH, Miller D, Winter VJ, Banfield MJ, Lee JH, Yoo SY, Henz SR, Brady RL, Weigel D. A divergent external loop confers antagonistic activity on floral regulators FT and TFL1. *EMBO J.* 2006;25:605–14.
4. Aki T, Shigyo M, Nakano R, Yoneyama T, Yanagisawa S. Nano scale proteomics revealed the presence of regulatory proteins including three FT-like proteins in Phloem and Xylem saps from rice. *Plant Cell Physiol.* 2008;49:767–90.
5. Bradley D, Ratcliffe O, Vincent C, Carpenter R, Coen E. Inflorescence commitment and architecture in *Arabidopsis*. *Science.* 1997;275:80–3.
6. Chardon F, Damerval C. Phylogenomic analysis of the PEBP gene family in cereals. *J Mol Evol.* 2005;61:579–90.
7. Conti L, Bradley D. TERMINAL FLOWER1 is a mobile signal controlling *Arabidopsis* architecture. *Plant Cell.* 2007;19:767–78.
8. Corbesier L, Vincent C, Jang S, Fornara F, Fan Q, Searle I, Giakountis A, Farrona S, Gissot L, Turnbull C, Coupland G. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science.* 2007;316:1030–3.

9. Daniel X, Sugano S, Tobin EM. CK2 phosphorylation of CCA1 is necessary for its circadian oscillator function in Arabidopsis. *Proc Natl Acad Sci USA*. 2004;101:3292–7.
10. Danilevskaya ON, Meng X, Hou ZL, Ananiev EV, Simmons CR. A genomic and expression compendium of the expanded PEBP gene family from maize. *Plant Physiol*. 2007;146:250–64.
11. Danilveskaya ON, Meng X, Selinger DA, Deschamps S, Hermon P, Vansant G, Gupta R, Ananiev EV, Muszynski MG. Involvement of the MADS-box gene ZMM4 in floral induction and inflorescence development in *Zea mays*. *Plant Physiol*. 2008. doi:10.1104/pp.107.115261.
12. Doi K, Izawa T, Fuse T, Yamanouchi U, Kubo T, Shimatani Z, Yano M, Yoshimura A. Ehd1, a B-type response regulator in rice, confers short-day promotion of flowering and controls FT-like gene expression independently of Hd1. *Genes Dev*. 2004;18:926–36.
13. Furutani I, Sukegawa S, Kyojuka J. Genome-wide analysis of spatial and temporal gene expression in rice panicle development. *Plant J*. 2006;46:503–11.
14. Giavalisco P, Kapitzka K, Kolasa A, Buhtz A, Kehr J. Towards the proteome of *Brassica napus* phloem sap. *Proteomics*. 2006;6:896–909.
15. Gu Q, Ferrándiz C, Yanofsky MF, Martienssen R. The FRUIT-FULL MADS-box gene mediates cell differentiation during *Arabidopsis* fruit development. *Development*. 1998;125:1509–17.
16. Hanzawa Y, Money T, Bradley D. A single amino acid converts a repressor to an activator of flowering. *Proc Natl Acad Sci USA*. 2005;102:7748–53.
17. Hayama R, Coupland G. The molecular basis of diversity in the photoperiodic flowering responses of *Arabidopsis* and rice. *Plant Physiol*. 2004;135:677–84.
18. Hayama R, Izawa T, Shimamoto K. Isolation of rice genes possibly involved in the photoperiodic control of flowering by a fluorescent differential display method. *Plant Cell Physiol*. 2002;43:494–504.
19. Hayama R, Yokoi S, Tamaki S, Yano M, Shimamoto K. Adaptation of photoperiodic control pathways produces short-day flowering in rice. *Nature*. 2003;422:719–22.
20. Hemming MN, Peacock WJ, Dennis ES, Trevaskis B. Low-temperature and daylength cues are integrated to regulate FLOWERING LOCUS T in barley. *Plant Physiol*. 2008;14:355–66.
21. Imaizumi T, Kay SA. Photoperiodic control of flowering: not only by coincidence. *Trends Plant Sci*. 2006;11:550–8.
22. Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA. FKF1F-BOX protein mediates cyclic degradation of a repressor of CONSTANS in *Arabidopsis*. *Science*. 2005;309:293–7.
23. Ishikawa R, Tamaki S, Yokoi S, Inagaki N, Shinomura T, Takano M, Shimamoto K. Suppression of the floral activator Hd3a is the principal cause of the night break effect in rice. *Plant Cell*. 2005;17:3326–36.
24. Izawa T. Daylength measurements by rice plants in photoperiodic short-day flowering. *Int Rev Cytol*. 2007a;256:191–222.
25. Izawa T. Adaptation of flowering-time by natural and artificial selection in *Arabidopsis* and rice. *J Exp Bot*. 2007b;58:3091–7.
26. Izawa T, Takahashi Y, Yano M. Comparative biology comes into bloom: genomic and genetic comparison of flowering pathways in rice and *Arabidopsis*. *Curr Opin Plant Biol*. 2003;6:113–20.
27. Izawa T, Oikawa T, Tokutomi S, Okuno K, Shimamoto K. Phytochromes confer the photoperiodic control of flowering in rice (a short-day plant). *Plant J*. 2000;22:391–9.
28. Izawa T, Oikawa T, Sugiyama N, Tanisaka T, Yano M, Shimamoto K. Phytochrome mediates the external light signal to repress FT orthologs in photoperiodic flowering of rice. *Genes Dev*. 2002;16:2006–20.
29. Jaeger KE, Wigge PA. FT protein acts as a long-range signal in *Arabidopsis*. *Curr Biol*. 2007;17:1050–4.
30. Jang S, Marchal V, Panigrahi KCS, Wenkel S, Soppe W, Deng XW, Valverde F, Coupland G. *Arabidopsis* COP1 shapes the temporal pattern of CO accumulation conferring a photoperiodic flowering response. *EMBO J*. 2008;27:1277–88.
31. Jeon JS, Lee S, Jung KH, Yang WS, Yi GH, Oh BG, An GH. Production of transgenic rice plants showing reduced heading date and plant height by ectopic expression of rice MADS-box genes. *Mol Breed*. 2000;6:581–92.
32. Kardailsky I, Shukla VK, Ahn JH, Dagenais N, Christensen SK, Nguyen JT, Chory J, Harrison MJ, Weigel D. Activation tagging of the floral inducer FT. *Science*. 1999;286:1962–5.
33. Kempin SA, Savidge B, Yanofsky MF. Molecular basis of the cauliflower phenotype in *Arabidopsis*. *Science*. 1995;267:522–5.
34. Kim SL, Lee SY, Kim HJ, Nam HG, An GH. OsMADS51 is a short-day flowering promoter that functions upstream of Ehd1, OsMADS14, and Hd3a(1[W][OA]). *Plant Physiol*. 2007a;145:1484–94.
35. Kim WY, Fujiwara S, Suh SS, Kim J, Kim Y, Han LQ, David K, Putterill J, Nam HG, Somers DE. ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature*. 2007b;449:356–60.
36. Kobayashi Y, Weigel D. Move on up, it's time for change—mobile signals controlling photoperiod-dependent flowering. *Genes Dev*. 2007;21:2371–84.
37. Kobayashi Y, Kaya H, Goto K, Iwabuchi M, Araki T. A pair of related genes with antagonistic roles in mediating flowering signals. *Science*. 1999;286:1960–2.
38. Kojima S, Takahashi Y, Kobayashi Y, Monna L, Sasaki T, Araki T, Yano M. Hd3a, a rice ortholog of the *Arabidopsis* FT gene, promotes transition to flowering downstream of Hd1 under short-day conditions. *Plant Cell Physiol*. 2002;43:1096–105.
39. Komiya R, Ikegami A, Tamaki S, Yokoi S, Shimamoto K. Hd3a and RFT1 are essential for flowering in rice. *Development*. 2008;135:767–74.
40. Kramer A, Feilner T, Possling A, Radchuk V, Weschke W, Burkle L, Kersten B. Identification of barley CK2 alpha targets by using the protein microarray technology. *Phytochemistry*. 2004;65:1777–84.
41. Kyojuka J, Kobayashi T, Morita M, Shimamoto K. Spatially and temporally regulated expression of rice MADS box genes with similarity to *Arabidopsis* class A, B and C genes. *Plant Cell Physiol*. 2000;41:710–8.
42. Laubinger S, Marchal V, Gentilhomme J, Wenkel S, Adrian J, Jang S, Kulajta C, Braun H, Coupland G, Hoecker U. *Arabidopsis* SPA proteins regulate photoperiodic flowering and interact with the floral inducer CONSTANS to regulate its stability. *Development*. 2006;133:3213–22.
43. Lee S, An G. Diversified mechanisms for regulating flowering time in a short-day plant rice. *J Plant Biol*. 2007;50:241–8.
44. Lee SY, Kim J, Han JJ, Han MJ, An GH. Functional analyses of the flowering time gene OsMADS50, the putative SUPPRESSOR OF OVEREXPRESSION OF CO 1/AGAMOUS-LIKE 20 (SOC1/AGL20) ortholog in rice. *Plant J*. 2004;38:754–64.
45. Li C, Dobcovsky J. Wheat FT protein regulates *VRN1* transcription through interactions with FDL2. *Plant J*. 2008. doi:10.1111/j.1365-3113X.2008.03526.x
46. Lin HX, Yamamoto T, Sasaki T, Yano M. Characterization and detection of epistatic interactions of 3 QTLs, Hd1, Hd2, and Hd3, controlling heading date in rice using nearly isogenic lines. *Theor Appl Genet*. 2000;101:1021–8.
47. Lin MK, Belanger H, Lee YJ, Varkonyi-Gasic E, Taoka KI, Miura E, Xoconostle-Cazares B, Gendler K, Jorgensene RA, Phinney B, Lough TJ, Lucas WJ. FLOWERING LOCUS T protein may act as the long-distance florigenic signal in the cucurbits. *Plant Cell*. 2007;19:1488–506.
48. Liu C, Chen H, Er HL, Soo HM, Kumar PP, Han JH, Liou YC, Yu H. Direct interaction of AGL24 and SOC1 integrates

- flowering signals in *Arabidopsis*. *Development*. 2008a;135:1481–91.
49. Liu LJ, Zhang YC, Li QH, Sang Y, Mao J, Lian HL, Wang L, Yan HQ. COP1-mediated ubiquitination of CONSTANS is implicated in cryptochrome regulation of flowering in *Arabidopsis*. *Plant Cell*. 2008b;20:292–306.
 50. Locke JCW, Kozma-Bognar L, Gould PD, Feher B, Kevei E, Nagy F, Turner MS, Hall A, Millar AJ. Experimental validation of a predicted feedback loop in the multi-oscillator clock of *Arabidopsis thaliana*. *Mol Syst Biol*. 2006;2:59.
 51. Mandel MA, Gustafson-Brown C, Savidge B, Yanofsky MF. Molecular characterization of the *Arabidopsis* floral homeotic gene *APETALA1*. *Nature*. 1992;360:273–7.
 52. Mathieu J, Warthmann N, Küttner F, Schmid M. Export of FT protein from phloem companion cells is sufficient for floral induction in *Arabidopsis*. *Curr Biol*. 2007;17:1055–60.
 53. Michaels SD, Himelblau E, Kim SY, Schomburg FM, Amasino RM. Integration of flowering signals in winter-annual *Arabidopsis*. *Plant Physiol*. 2005;137:149–56.
 54. Mizoguchi T, Wright L, Fujiwara S, Cremer F, Lee K, Onouchi H, Mouradov A, Fowler S, Kamada H, Putterill J, Coupland G. Distinct roles of GIGANTEA in promoting flowering and regulating circadian rhythms in *Arabidopsis*. *Plant Cell*. 2005;17:2255–70.
 55. Monna L, Lin HX, Kojima S, Sasaki T, Yano M. Genetic dissection of a genomic region for a quantitative trait locus, Hd3, into two loci, Hd3a and Hd3b, controlling heading date in rice. *Theor Appl Genet*. 2002;104:772–8.
 56. Murai K, Miyamae M, Kato H, Takumi S, Ogiwara Y. WAP1, a wheat *APETALA1* homolog, plays a central role in the phase transition from vegetative to reproductive growth. *Plant Cell Physiol*. 2003;44:1255–65.
 57. Murakami M, Tago Y, Yamashino T, Mizuno T. Comparative overviews of clock-associated genes of *Arabidopsis thaliana* and *Oryza sativa*. *Plant Cell Physiol*. 2007;48:110–21.
 58. Muszynski MG, Dam T, Li B, Shirbroun DM, Hou ZL, Bruggemann E, Archibald R, Ananiev EV, Danilevskaya ON. Delayed flowering1 encodes a basic leucine zipper protein that mediates floral inductive signals at the shoot apex in maize. *Plant Physiol*. 2006;142:1523–36.
 59. Nakagawa M, Shimamoto K, Kyozuka J. Overexpression of RCN1 and RCN2, rice TERMINAL FLOWER 1/CENTRORADIALIS homologs, confers delay of phase transition and altered panicle morphology in rice. *Plant J*. 2002;29:743–50.
 60. Nijhawan A, Jain M, Tyagi AK, Khurana JP. Genomic survey and gene expression analysis of the basic leucine zipper transcription factor family in rice. *Plant Physiol*. 2008;146:333–50.
 61. Niwa Y, Ito S, Nakamichi N, Mizoguchi T, Niinuma K, Yamashino T, Mizuno T. Genetic linkages of the circadian clock-associated genes, TOC1, CCA1 and LHY, in the photoperiodic control of flowering time in *Arabidopsis thaliana*. *Plant Cell Physiol*. 2007;48:925–37.
 62. Page T, Macknight R, Yang CH, Dean C. Genetic interactions of the *Arabidopsis* flowering time gene FCA, with genes regulating floral initiation. *Plant J*. 1999;17:231–9.
 63. Portoles S, Mas P. Altered oscillator function affects clock resonance and is responsible for the reduced day-length sensitivity of CKB4 overexpressing plants. *Plant J*. 2007;51:966–77.
 64. Riera M, Figueras M, Lopez C, Goday A, Pages M. Protein kinase CK2 modulates developmental functions of the abscisic acid responsive protein Rab17 from maize. *Proc Natl Acad Sci USA*. 2004;101:9879–84.
 65. Samach A, Onouchi H, Gold SE, Ditta GS, Schwarz-Sommer Z, Yanofsky MF, Coupland G. Distinct roles of CONSTANS target genes in reproductive development of *Arabidopsis*. *Science*. 2000;288:1613–6.
 66. Sawa M, Nusinow DA, Kay SA, Imaizumi T. FKF1 and GIGANTEA complex formation is required for day-length measurement in *Arabidopsis*. *Science*. 2007;318:261–5.
 67. Shitsukawa N, Ikari C, Shimada S, Kitagawa S, SakaMoto K, Saito H, Ryuto H, Fukunishi N, Abe T, Takumi S, Nasuda S, Murai K. The einkorn wheat (*Triticum monococcum*) mutant, maintained vegetative phase, is caused by a deletion in the VRN1 gene. *Genes Genet Syst*. 2007;82:167–70.
 68. Stemmer C, Schwander A, Bauw G, Fojan P, Grasser KD. Protein kinase CK2 differentially phosphorylates maize chromosomal high mobility group B (HMGB) proteins modulating their stability and DNA interactions. *J Biol Chem*. 2002;277:1092–8.
 69. Sugano S, Andronis C, Green RM, Wang ZY, Tobin EM. Protein kinase CK2 interacts with and phosphorylates the *Arabidopsis* circadian clock-associated 1 protein. *Proc Natl Acad Sci USA*. 1998;95:11020–5.
 70. Sugano S, Andronis C, Ong MS, Green RM, Tobin EM. The protein kinase CK2 is involved in regulation of Circadian rhythms in *Arabidopsis*. *Proc Natl Acad Sci USA*. 1999;96:12362–6.
 71. Takada S, Goto K. TERMINAL FLOWER2, an *Arabidopsis* homolog of HETEROCHROMATIN PROTEIN1, counteracts the activation of FLOWERING LOCUS T by CONSTANS in the vascular tissues of leaves to regulate flowering time. *Plant Cell*. 2003;15:2856–65.
 72. Takahashi Y, Shomura A, Sasaki T, Yano M. Hd6, a rice quantitative trait locus involved in photoperiod sensitivity, encodes the alpha subunit of protein kinase CK2. *Proc Natl Acad Sci USA*. 2001;98:7922–7.
 73. Takano M, Inagaki N, Xie XZ, Yuzurihara N, Hihara F, Ishizuka T, Yano M, Nishimura M, Miyao A, Hirochika H, Shinomura T. Distinct and cooperative functions of phytochromes A, B, and C in the control of deetiolation and flowering in rice. *Plant Cell*. 2005;17:3311–25.
 74. Tamaki S, Matsuo S, Wong HL, Yokoi S, Shimamoto K. Hd3a protein is a mobile flowering signal in rice. *Science*. 2007;316:1033–6.
 75. Teper-Bamnolker P, Samach A. The flowering integrator FT regulates SEPALLATA3 and FRUITFULL accumulation in *Arabidopsis* leaves. *Plant Cell*. 2005;17:2661–75.
 76. Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D. Integration of spatial and temporal information during floral induction in *Arabidopsis*. *Science*. 2005;309:1056–9.
 77. Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q. Natural variation in *Ghd7* is an important regulator of heading date and yield potential in rice. *Nat Genet*. 2008;40:761–7.
 78. Yamaguchi A, Kobayashi Y, Goto K, Abe M, Araki T. TWIN SISTER OF FT (TSF) acts as a floral pathway integrator redundantly with FT. *Plant Cell Physiol*. 2005;46:1175–89.
 79. Yamamoto T, Lin HX, Sasaki T, Yano M. Identification of heading date quantitative trait locus Hd6 and characterization of its epistatic interactions with Hd2 in rice using advanced backcross progeny. *Genetics*. 2000;154:885–91.
 80. Yamamoto T, Kuboki Y, Lin SY, Sasaki T, Yano M. Fine mapping of quantitative trait loci Hd-1, Hd-2 and Hd-3, controlling heading date of rice, as single Mendelian factors. *Theor Appl Genet*. 1998;97:37–44.
 81. Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J. Positional cloning of the wheat vernalization gene VRN1. *Proc Natl Acad Sci USA*. 2003;100:6263–8.
 82. Yan L, Fu D, Li C, Blechl A, Tranquilli G, Bonafede M, Sanchez A, Valarik M, Yasuda S, Dubcovsky J. The wheat and barley vernalization gene VRN3 is an orthologue of FT. *Proc Natl Acad Sci USA*. 2006;103:19581–6.

83. Yano M, Kojima S, Takahashi Y, Lin HX, Sasaki T. Genetic control of flowering time in rice, a short-day plant. *Plant Physiol.* 2001;127:1425–9.
84. Yano M, Katayose Y, Ashikari M, Yamanouchi U, Monna L, Fuse T, Baba T, Yamamoto K, Umehara Y, Nagamura Y, Sasaki T. Hd1, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the *Arabidopsis* flowering time gene *CONSTANS*. *Plant Cell.* 2000;12:2473–83.
85. Zeevaart JAD. Florigen coming of age after 70 years. *Plant Cell.* 2006;18:1783–9.
86. Zeilinger MN, Farre EM, Taylor SR, Kay SA, Doyle FJ. A novel computational model of the circadian clock in *Arabidopsis* that incorporates PRR7 and PRR9. *Mol Syst Biol.* 2006;2:58.
87. Zhang SH, Hu WJ, Wang LP, Lin CF, Cong B, Sun CR, Luo D. TFL1/CEN-like genes control intercalary meristem activity and phase transition in rice. *Plant Science.* 2005;168:1393–408.